




Breast Screening Options

This table outlines the differences between mammography, thermography, and ultrasound.

<p>Mammography</p> 	<p>Thermography</p> 	<p>Ultrasound</p> 
<p>Uses X-rays to produce an image that is a shadow of dense structures. Suspicious areas need to be dense enough to be seen.</p>	<p>Uses infrared sensors to detect heat and increased vascularity (angiogenesis) as the byproduct of biochemical reactions. The heat is compiled into an image for computerized analysis.</p>	<p>High frequency sound waves are bounced off the breast tissue and collected as an echo to produce an image.</p>
<p>Structural imaging. Ability to locate the area of suspicious tissue.</p>	<p>Functional imaging. Detects physiologic changes. Cannot locate the exact area of suspicion inside the breast.</p>	<p>Structural imaging. Ability to locate the area of suspicious tissue.</p>
<p>Early detection method.</p>	<p>Early detection method. Used as an adjunctive imaging test.</p>	<p>Low spatial resolution (cannot see fine detail). Good at distinguishing solid masses from fluid filled cysts. Used as an adjunctive imaging test.</p>
<p>Findings increase suspicion. Cannot diagnose cancer.</p>	<p>Findings increase suspicion. Cannot diagnose cancer.</p>	<p>Findings increase suspicion. Cannot diagnose cancer.</p>
<p>Can detect tumors in the pre- invasive stage.</p>	<p>May provide the first signal that a problem is developing.</p>	<p>Ability to detect some cancers missed by mammography.</p>
<p>No comparison.</p>	<p>A positive infrared image represents the highest known risk factor for the existence of or future development of breast cancer – 10 times more significant than any family history of the disease.</p>	<p>No comparison.</p>
<p>Average 80% Sensitivity (20% of cancers missed), in women over age 50. Sensitivity drops to 60% (40% of cancers missed) in women under age 50.</p>	<p>Average 90% Sensitivity (10% of cancers missed) in all age groups.</p>	<p>Average 83% Sensitivity (17% of cancers missed) in all age groups.</p>
<p>Hormone use decreases sensitivity.</p>	<p>No known effect.</p>	<p>No known effect.</p>
<p>Large, dense, and fibrocystic breasts cause reading difficulties.</p>	<p>No effect.</p>	<p>No known effect.</p>
<p>In most women, the medial upper triangle, peripheral areas next to the chest wall, and the inframammary sulcus cannot be visualized.</p>	<p>All areas visualized.</p>	<p>All areas visualized.</p>

A biopsy is the only test that can determine if a suspected tissue area is cancerous.